

EmerAid® IntensiveCare HDN™ Kidney

Critical Care Nutrition for Felines with Chronic Kidney Disease



Highest kcal/ml for Feline Kidney Disease: **2.23kcal/ml**

- Highly digestible essential amino acids like L-carnitine to support lean muscle mass and combat cachexia
- Controlled phosphorous levels help delay the progression of CKD
- Omega 3 fatty acids – EPA & DHA – aid in slowing CKD progression
- Antioxidants to minimize oxidative stress
- Semi-elemental diet = Fastest nutrient absorption
- Easy to administer through a tube as small as 5Fr
- 30-day shelf life after opening



Manufactured by: EmerAid LLC, A Division of Lafeber 24981 N 1400 East Road, Cornell, IL 61319
1.815.466.8040 customerservice@EmerAid.com EmerAidVet.com  MADE IN THE USA

EmerAid® Intensive Care HDN™ Kidney

Feline IC HDN Kidney kcal/ml Feeding Chart


Feline		70 x BW(kg) ^{0.75} RER (kcal/day)	1.45 kcal/ml 5Fr - 8Fr Tube		1.81 kcal/ml 8Fr - Larger Tube		2.23 kcal/ml Syringe				
Weight KG	Weight LB		Total ml/day	TID	QID	Total ml/day	TID	QID	Total ml/day	TID	QID
1	2.2	70	48	16	12	39	14	10	31	10	8
2	4.4	118	81	27	20	65	22	16	53	18	13
3	6.6	160	110	37	28	88	29	22	72	24	18
4	9	198	137	46	34	109	36	27	89	30	22
5	11	234	161	54	40	129	43	32	105	35	26
6	13	268	185	62	46	148	49	37	120	40	30
8	18	333	230	77	58	184	61	46	149	50	37
10	22	394	272	91	68	218	73	55	177	59	44
12	26	451	311	104	78	249	83	62	202	67	51
14	31	507	350	117	88	280	93	70	227	76	57

Feline IC HDN Kidney Mixing Chart

EmerAid Powder (parts)	Water* (parts)	Feeding Route	Kcal/ml**
3	1	HDN Bites	2.81
2	1	Syringe	2.23
1.5	1	10Fr and Larger Tube	1.81
1	1	5Fr - 8Fr Tube	1.45

Measure out
EmerAid
powder

Make fresh
each time



*Mix with hot water measuring 122°F (50°C). **Metabolizable energy when made as directed.

Standard Mixing Directions: Measure out EmerAid according to the proportions needed. Mix EmerAid using the scoop provided. Mix with hot water measuring 122°F (50°C). Mix well. Use a syringe to draw up the mixture. When finished, flush and clean the feeding syringe. Make fresh for each feeding. Discard remaining food after each feeding.

Feeding frequency and volume should be adjusted based upon patient response and under the supervision of a veterinarian.

Use within 30 days of opening.

29.02 kcal/scoop



Manufactured by: EmerAid LLC, A Division of Lafeber 24981 N 1400 East Road, Cornell, IL 61319
1.815.466.8040 customerservice@EmerAid.com EmerAidVet.com MADE IN THE USA